


PATENT  
ATTORNEY DOCKET NO. 50025/003003

Certificate of Mailing	
Date of Deposit <u>June 5, 2001</u>	Label Number <u>EL509219168US</u>
I hereby certify under 37 C.F.R. § 1.10 that this correspondence is being deposited with the United States Postal Service as " <b>Express Mail Post Office to Addressee</b> " with sufficient postage on the date indicated above and is addressed to BOX PATENT APPLICATION, Assistant Commissioner for Patents, Washington, D.C. 20231.	
Guy Beardsley	
Printed name of person mailing correspondence	Signature of person mailing correspondence

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant:	Victor J. Dzau <i>et al.</i>	Art Unit:	Not Yet Assigned
Serial No.:	Not Yet Assigned	Examiner:	Not Yet Assigned
Filed:	June 5, 2001		
Title:	Therapeutic Use of Cis-Element Decoys <i>In Vivo</i>		

Box Patent Application  
Assistant Commissioner for Patents  
Washington, D.C. 20231

PRELIMINARY AMENDMENT AND REMARKS

Applicants submit the following Amendment and Remarks for consideration prior to examination of the above-captioned patent application.

09375305-060501  
T05090"060501

## AMENDMENT

Kindly amend the application as follows.

### In the Claims:

Cancel claims 1-12, without prejudice, and add the following new claims 13-27.

13. (New) A method for preventing or treating an NFκB -associated disease or condition in a mammal, said method comprising introducing an NFκB decoy into a cell of said mammal.

14. (New) The method of claim 13, wherein said mammal is a human.

15. (New) The method of claim 13, wherein said decoy is introduced into said cell *ex vivo*.

16. (New) The method of claim 13, wherein said decoy is introduced into said cell *in vivo*.

17. (New) The method of claim 13, wherein said decoy comprises a polynucleotide.

18. (New) The method of claim 17, wherein said decoy comprises dsDNA.

19. (New) The method of claim 14, wherein said human has or is at risk of developing ischemic reperfusion injury.

20. (New) The method of claim 14, wherein said human has or is at risk of developing myocardial infarction.

21. (New) The method of claim 14, wherein said human has or is at risk of developing inflammation.

22. (New) The method of claim 21, wherein said human has or is at risk of developing glomerulonephritis.

23. (New) The method of claim 21, wherein said human has or is at risk of developing dermatitis.

24. (New) The method of claim 14, wherein said human has or is at risk of developing immune disease.

25. (New) The method of claim 14, wherein said human has or is at risk of developing transplant rejection.



## REMARKS

The amendment set forth above adds to the application independent claim 13, as well as several claims that depend from claim 13. New claim 13 is drawn to a method of preventing or treating an NFκB-associated disease or condition in a mammal, in which an NFκB decoy is introduced into the cell. New claim 13 is similar to claim 8 of the parent application, U.S. Serial No. 08/524,206, except that claim 13 is narrower in that it specifies the use of a decoy to a particular transcription factor, NFκB.

Claim 8 was not rejected over prior art in the parent application. Moreover, the operability of the subject matter of claim 13, employing an NFκB decoy, is shown in the enclosed papers. Sawa et al., Circulation 96 (Suppl. II); II-280-II-285, 1997 (a copy is enclosed), for example, describes a study in which NFκB decoys were used to treat ischemic reperfusion injury in the rat myocardium. NFκB decoys were introduced into rat hearts by coronary infusion. Ischemic reperfusion injury in rats that were treated with the NFκB decoys was attenuated as compared to such injury in control rats, which were treated with scrambled decoys. NFκB decoy-treated rats also showed lower levels of neutrophil adherence to endothelial cells and decreased levels of IL-8, as compared to controls, further showing the efficacy of NFκB decoys in treating ischemic reperfusion injury.

In a second paper, Tomita et al., Arthritis & Rheumatism 42(12):2532-2542, 1999 (enclosed), a study is described in which NFκB decoys were used to treat inflammatory arthritis in rats. Briefly, NFκB decoys were introduced into the bilateral hind ankle joints

of rats with collagen-induced arthritis (CIA) by intraarticular injection. This treatment resulted in decreased severity of hind-paw swelling, suppression of joint destruction, and decreased production of IL-1 and TNF- $\alpha$  in the synovium of arthritic joints in NF $\kappa$ B-treated rats, as compared to controls, which were treated with a scrambled oligonucleotide.

Decoys were also shown to have effect in treating inflammation in a paper by D'Acquisto et al., *Gene Ther.* 7(20):1731-1737, 2000 (enclosed). In this study, local injection of NF $\kappa$ B decoys was shown to reduce inhibit edema formation induced by carrageenin in hind paws of rats.

NF $\kappa$ B decoys were shown to suppress experimental crescentic glomerulonephritis in a third paper, Tomita et al., *J. Am Soc. Nephrol.* 11(7):1244-1252, 2000 (enclosed). In this study, the decoys were introduced into the left kidneys of rats in which glomerulonephritis had been induced. This treatment resulted in substantial inhibition of disease, as shown by reduced proteinuria, histologic damage, leukocyte infiltration, and cytokine and leukocyte adhesion molecule expression, as compared to scrambled oligonucleotide-treated controls. Similar results were reported by Tomita et al., *Gene Ther.* 7(15):1326-1332, 2000 (enclosed).

In a paper by Khaled et al., *Clin. Immunol. Immunopathol.* (2):170-179, 1998 (enclosed), a study is reported in which NF $\kappa$ B decoys were shown to have an effect on immunity. In particular, the decoys were shown to inhibit cytokine production in splenocytes from an autoimmune mouse strain. Decoys were further shown to inhibit

tumor cell growth, *in vitro* and *in vivo*, in a paper by Sharma et al., Anticancer Res. 16(2):589-596, 1996 (enclosed). In particular, this review describes studies that showed that a decoy that is specific for the *RelA* subunit of NFκB blocked tumor cell growth in soft agar, as well as inhibited tumorigenicity in an *in vivo* tumor model.

The data described in each of the papers discussed above show that NFκB decoys are effective agents in the treatment of NFκB-associated diseases and conditions, as is specified in new claims 13-27.

### CONCLUSION

Although no charges are believed to be due, if there are any charges or any credits, please apply them to Deposit Account No. 03-2095.

Respectfully submitted,

Date:

June 4, 2001

Susan M. Michaud  
Susan M. Michaud, Ph.D.  
Reg. No. 42,885

Clark & Elbing LLP  
176 Federal Street  
Boston, MA 02110  
Telephone: 617-428-0200  
Facsimile: 617-428-7045  
50025.003003 preliminary amendment msword version.doc



21559

PATENT TRADEMARK OFFICE